



Autotrophic nitrogen assimilation and carbon capture for microbial protein production by a novel enrichment of hydrogen-oxidizing bacteria



Silvio Matassa ^{a, b}, Willy Verstraete ^{a, b}, Ilje Pikaar ^c, Nico Boon ^{a, *}

^a Center for Microbial Ecology and Technology (CMET), Ghent University, Coupure Links 653, 9000, Gent, Belgium

^b Avecom NV, Industrieweg 122P, 9032, Wondelgem, Belgium

^c School of Civil Engineering, The University of Queensland, Brisbane, QLD, 4072, Australia

ARTICLE INFO

Article history:

Received 24 March 2016

Received in revised form

20 May 2016

Accepted 24 May 2016

Available online 26 May 2016

Keywords:

Resource up-cycling

Nitrogen assimilation

Carbon capture

Microbial protein

Hydrogen-oxidizing bacteria

Sulfuricurvum spp.

ABSTRACT

Domestic used water treatment systems are currently predominantly based on conventional resource inefficient treatment processes. While resource recovery is gaining momentum it lacks high value end-products which can be efficiently marketed. Microbial protein production offers a valid and promising alternative by upgrading low value recovered resources into high quality feed and also food. In the present study, we evaluated the potential of hydrogen-oxidizing bacteria to upgrade ammonium and carbon dioxide under autotrophic growth conditions. The enrichment of a generic microbial community and the implementation of different culture conditions (sequenced batch resp. continuous reactor) revealed surprising features. At low selection pressure (i.e. under sequenced batch culture at high solid retention time), a very diverse microbiome with an important presence of predatory *Bdellovibrio* spp. was observed. The microbial culture which evolved under high rate selection pressure (i.e. dilution rate $D = 0.1 \text{ h}^{-1}$) under continuous reactor conditions was dominated by *Sulfuricurvum* spp. and a highly stable and efficient process in terms of N and C uptake, biomass yield and volumetric productivity was attained. Under continuous culture conditions the maximum yield obtained was 0.29 g cell dry weight per gram chemical oxygen demand equivalent of hydrogen, whereas the maximum volumetric loading rate peaked 0.41 g cell dry weight per litre per hour at a protein content of 71%. Finally, the microbial protein produced was of high nutritive quality in terms of essential amino acids content and can be a suitable substitute for conventional feed sources such as fishmeal or soybean meal.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

Primary producers - autotrophic microorganisms - are essential for carbon and nutrients cycling. While fixing inorganic CO_2 into organic biomass they recycle nutrients (N and P) and provide food for higher life forms (Elser et al., 2000). Primary producers such as algae and autotrophic bacteria can serve as alternative protein source in the form of microbial protein (MP) for livestock but also for human consumption (Anupama and Ravindra, 2000; Verstraete, 2015; Walsh et al., 2015). Besides protein, microbes can also accumulate considerable amounts of biocompatible prebiotics such as PHB (Defoirdt et al., 2007), thereby enhancing the nutritional value

of the microbial biomass.

After being extensively studied in the past, mainly as means to upgrade fossil fuel (e.g. paraffin, natural gas) to protein supplements (Westlake, 1986), the use of bacteria for microbial protein (MP) production has nowadays re-gained significant interest (Aas et al., 2006; Marit Berge et al., 2005) with natural gas based MP production entering the market economy (Strong et al., 2015). Innovative approaches implementing bacteria to produce MP within the context of resource recovery from used water have also been recently proposed (Lee et al., 2015; Liu et al., 2016; Matassa et al., 2015a). Indeed, the production of MP can allow the up-cycling of nitrogen and carbon dioxide recovered from used water streams, converting them into protein-rich feed and food substances. Different physico-chemical techniques can be implemented in the recovery of N and C substrates. Air stripping or pervaporative processes can recover N from concentrated streams

* Corresponding author.

E-mail address: Nico.Boon@UGent.be (N. Boon).

such as anaerobic digestate, whereas pressure swing adsorption (PSA) can concentrate CO₂ from biogas, thus providing the building blocks which are at the base of MP biosynthesis.

Among the various metabolic pathways suitable for MP production, including both eukaryotic and prokaryotic microorganisms (Anupama and Ravindra, 2000), autotrophic hydrogen-oxidizing bacteria (HOB) constitute a special and thus far unexplored metabolic niche with potential for novel applications in resource recovery and upgrade. Even if ubiquitous, autotrophic HOB have only received limited attention, with previous studies focusing on the use of axenic cultures comprising bacteria such as *Alcaligenes eutrophus*, *Ralstonia eutropha*, *Seliberia carboxydohydrogena* (Ishizaki and Tanaka, 1990; Repaske and Mayer, 1976; Volova and Barashkov, 2010). The metabolic features of autotrophic HOB allow them to grow on hydrogen (electron donor) and oxygen (electron acceptor) while fixing carbon dioxide into cell material and assimilating nitrogen into high quality protein (Parkin and Sargent, 2012; Pohlmann et al., 2006). MP produced by autotrophic HOB is characterized by all the essential amino acids, having an amino acid profile closer to high-quality animal protein rather than to vegetable protein (Volova and Barashkov, 2010). Given this interesting feature, autotrophic HOB were already proposed as possible protein source within biological life support systems for space missions (Bartsev et al., 1996), as well as for human and animal nutrition (Volova and Barashkov, 2010).

An attractive characteristic of MP production with autotrophic HOB is the possibility to exploit the increasing potential of renewable energy generation. A clear example is the use of hydrogen gas produced from water electrolysis, powered by e.g. wind or solar energy, or also from biomass gasification (Ni et al., 2006). Recently, biomethane has also been proposed as possible renewable feedstock for hydrogen production by means of a combined heat, hydrogen and power generation unit (CHHP) (Agll et al., 2013; Hamad et al., 2014). The possibility to implement such technologies on-site and produce hydrogen on demand might enable the direct up-cycling of mineral nitrogen and carbon dioxide recovered from wastewater treatment plants, as previously mentioned. Moreover, upcoming technological developments and the decrease of hydrogen prices (Ball and Weeda, 2015) justify further research efforts towards the application of autotrophic HOB within resource recovery and up-cycling.

In the present study, we aimed to experimentally determine the feasibility of nitrogen and carbon upgrade into MP by means of a microbial community enriched in HOB using a lab-scale gas. Along the experimental investigation different culture conditions were imposed to the enriched HOB culture (i.e. sequenced batch and continuous). This was done in order to establish how the microbial community was shaped by the process conditions and how this affected the overall biological performance of the system, aiming at maximizing MP production (i.e. biomass yield and volumetric productivities). Nitrogen under the form of ammonium salt and gaseous CO₂ represented the N and C substrates needed for the production of MP protein by means of autotrophic HOB. The study started with the enrichment of a generic aerobic microbial mixed culture with autotrophic HOB under sequencing batch reactor operations. Consequently, the enriched mixed community was cultured in a continuous reactor configuration, resulting in the ongoing evolvement of a highly specific bacterial culture dominated by the genus *Sulfuricurvum*. The efficiency of the process in terms of gas utilization and by-product formation was monitored along the time course of the selective enrichment process. The microbial community analyses of the HOB microbiome under batch and continuous culture systems allowed delineating the evolution of the mixed bacterial community towards a quasi-monoculture dominated by *Sulfuricurvum* spp. Finally, the MP produced was

characterized in terms of crude protein content and amino acid profile in order to assess its nutritional value.

2. Material and methods

2.1. Enrichment of hydrogen-oxidizing bacteria

Aerobic sludge from a local food (potatoes) processing plant (Nazareth, Belgium) was used as an initial mixed culture for the enrichment of autotrophic HOB community. The enrichment was carried out in a 1 L gas fermentor. The fermentor was connected to 3 gas bags supplying a gas mixture composed by H₂/O₂/CO₂ with the following composition: 65/20/15 (vol/vol). Prior to use, each gas bag was flushed with Alphagaz 2-grade H₂, O₂ and CO₂ gases (Air Liquide, Belgium). The gaseous H₂/O₂/CO₂ atmosphere was constantly recirculated between the culture vessels and the gas bags by means of a peristaltic pump adapted to gas recirculation (Sci-Q 300, Watson Marlow, Belgium). The reactor was placed in a 28 °C temperature controlled room and shaken at 150 rpm. A volume of 500 mL of mineral media inoculated with 10% of inoculum was used at start. The mineral medium was prepared accordingly to Yu et al. (2013) for HOB isolation and culturing. The growth of HOB was followed by monitoring the increase of cell dry weight (CDW) over the course of the experimental run. When ammonium nitrogen was depleted, 50 mL of bacterial culture was withdrawn and diluted into 450 mL of fresh medium to restart the enrichment. After a stable and reproducible growth was attained in terms of CDW concentrations (2–3 g CDW/L before medium replenishment), the culture was considered enriched and used to start the experimental phase in the final reactor setup.

2.2. Reactor operations and controls

A completely stirred tank reactor (CSTR) (Biostat A plus, Sartorius, Belgium) was used during batch as well as continuous experiments. The 5 L glass vessel, with a working volume of 3 L, was stirred at 700 rpm with a 3-blade segment impeller to ensure completely mixed conditions. Hydrogen gas was produced on site by means of a lab-grade hydrogen generator (Alphagaz™ Flo H2, Air Liquide, Belgium), while CO₂ from gas bottles was of the same grade of the one used during the initial enrichment of the culture. Compressed air was used to provide the oxygen. Gases were fed separately by means of 3 micro-spargers (Sartorius, Belgium) submerged in the reactor. Gas flows were monitored using gas rotameters (Omega, USA) and kept at H₂: 120 mL/min; CO₂: 25 mL/min; Air: 180 mL/min. The gas collected in the headspace was constantly recirculated by means of a peristaltic pump adapted to gas recirculation (Sci-Q 300, Watson Marlow, Belgium) using a fourth micro-sparger. Utilized gas by the bacteria, was bubbled through an external water lock (imposing an overpressure of 20 mbar) and subsequently vented to the atmosphere by means of a fume hood. Temperature and pH were automatically controlled and kept at 35 ± 1 °C and 6.7, respectively.

2.3. Sequencing batch and continuous reactor culture systems

Sequencing batch reactor (SBR) tests were started by transferring 300 mL of fully grown bacterial culture into 2.7 L of fresh mineral medium, allowing an initial cell dry weight Cell Dry Weight (CDW) concentration of 300–500 mg CDW/L. Each sequencing batch test was allowed to evolve for an average of 5–6 days before transferring the culture into fresh medium, corresponding to a solid retention time (SRT) of 6 ± 0.5 days. Additional NH₄Cl was added to the standard mineral medium composition in order to achieve initial NH₄⁺-N concentration of 1.2 g/L, and simulate higher N

concentrations obtainable with recovery techniques such as air stripping or pervaporative systems. The sequencing batch culture was monitored along a period of 5 months.

Continuous reactor (CR) operations were set by supplying fresh media with a diaphragm pump (Qdos, Watson Marlow, Belgium), totaling a flow of 7.2 L/day. In the same way, 7.2 L/day of cell culture were constantly withdrawn from the CSTR reactor by means of a similar pump. As for the sequenced batch experiment, additional NH₄Cl was added to the standard mineral medium composition in order to achieve initial NH₄⁺-N concentration of 0.5 g/L. The complete absence of biomass recirculation set hydraulic and (SRT) of 10 h. Under these continuous reactor conditions (chemostat), only bacteria with a specific growth rate “μ” equal or higher than the dilution rate $D = 0.1 \text{ h}^{-1}$ could avoid being washed-out from the biological system. The continuous system was operated uninteruptedly for 3 months.

2.4. Analytical methods

NH₄⁺-N concentrations were determined by means of cuvette

$$\text{Gas conversion efficiency}(\%) = \frac{\text{Gas inlet (mol/min)} - \text{Gas outlet (mol/min)}}{\text{Gas inlet (mol/min)}} \times 100 \quad (1)$$

tests (Hach Lange, range 0–47 mg NH₄⁺-N/L). Cell Dry Weight (CDW) was measured in duplicate after water was evaporated at 105 °C for 24 h. Prior to analysis, the samples were centrifuged at 12500 g for 10 min for three times, each time re-suspending the biomass pellet in demineralized water. Gas samples collected from the reactor headspace were analyzed with a Compact GC (Global Analyser Solutions, Breda, The Netherlands), equipped with a Molsieve 5A pre-column and Porabond column (O₂, H₂ and N₂) and a Rt-Q-bond pre-column and column (CO₂). Concentrations of gases were determined by means of a thermal conductivity detector.

2.5. Analysis and characterization of microbial protein

Kjeldahl nitrogen content of the microbial biomass was analyzed according to Standard methods (APHA et al., 1992). Organic nitrogen was determined as the difference between Kjeldahl nitrogen and ammonium nitrogen. The final protein content of CDW was obtained by multiplying the obtained value by applying a conversion factor of 6.25 as done in previous studies (Saloväänänen and Koivistoinen, 1996). The dietary amino acids composition of the microbial biomass was determined by an external accredited commercial laboratory (Eurofins Denmark A/S, Denmark).

2.6. Microbial community analysis

Liquid samples for total DNA extraction were centrifuged for 10 min at 10,000 RPM. Subsequently, the supernatant was removed and biomass pellet was stored immediately at –20 °C until further analysis following a protocol adapted from Vilchez-Vargas et al. (2013). Cells were lysed with 1 mL lysis buffer (100 mM Tris/HCl pH 8.0, 100 mM EDTA pH 8, 100 mM NaCl, 1% (m/v) polyvinylpyrrolidone and 2% (m/v) sodium dodecyl sulphate) and 200 mg glass beads (0.11 mm, Sartorius) in a FastPrepR - 96 instrument (MP Biomedicals, Santa Ana, USA) for two times 40 s

(1600 rpm). After removing glass beads by centrifugation (5 min at 10,000 RPM), DNA was extracted from supernatant following a phenol–chloroform extraction. DNA was precipitated with 1 vol ice-cold isopropyl alcohol and 0.1 vol 3 M sodium acetate for at least 1 h at –20 °C. After removal of isopropyl alcohol by centrifugation (30 min, maximum speed), the DNA pellet was dried and re-suspended in 100 μL 1 × TE (10 mM Tris, 1 mM EDTA) buffer. After finishing the extraction protocol, the DNA samples were immediately stored at –20 °C until further processing. Quality of DNA samples was analyzed by 1% (w/v) agarose (Life technologies, Madrid, Spain) gel electrophoresis. The PCR amplicons were purified with the innuPREP PCR pure kit (Analytik Jena, Jena, Germany), and sequenced with the primers used for PCR. 16s rRNA Illumina and Sanger sequencing analyses were conducted for each sample in triplicate by external commercial laboratories (Analytik Jena, Jena, Germany).

2.7. Calculations

The gas conversion efficiency was calculated as:

$$Y_{\text{H}_2} \left(\frac{\text{gCDW}}{\text{gH}_2 - \text{COD}} \right) = \frac{\text{CDW(g/L)}}{\text{H}_2 \text{ gas uptake(mol)} \times 16(\text{gCOD/mol}) \times \text{Liquid volume(L)}} \quad (2)$$

The biomass yield on carbon dioxide is calculated as:

$$Y_{\text{CO}_2} \left(\frac{\text{gCDW} - C}{\text{gCO}_2 - C} \right) = \frac{\text{CDW(g/L)} \times 0.5(\text{gC/gCDW})}{\text{CO}_2 \text{ gas uptake(mol)} \times 12(\text{gC/mol}) \times \text{Liquid volume(L)}} \quad (3)$$

The mineral nitrogen upgrade efficiency is calculated as:

$$\text{N upgrade efficiency}(\%) = \frac{\text{NH}_4 - \text{N in(g/L)} - X - \text{N out(g/L)}}{\text{NH}_4 - \text{N in}} \times 100 \quad (4)$$

Where NH₄-N in indicates the amount of NH₄-N fed to the reactor, respectively to the SBR and the CR systems and X-N out indicates the amount of dissolved nitrogen under the form of NH₄⁺, NO₂⁻ or NO₃⁻ at in at the end of each SBR test and in the effluent of the CR system.

3. Results

3.1. Sequencing batch and continuous reactor performances

The enriched HOB culture was first cultivated under sequencing batch reactor (SBR) conditions, with a SRT of 6 ± 0.5 days. The same experimental setup was then adapted to grow the HOB culture under continuous reactor (CR) configuration, imposing a SRT of

10 h. The main parameters analyzed both under SBR and CR configurations were: volumetric productivities (g CDW/L·h), biomass yields on hydrogen (g CDW/g COD-H₂) and carbon dioxide (g CDW-C/g CO₂-C) and hydrogen gas conversion efficiencies (%), as shown in Table 1.

The average values for each individual SBR test reported in Table 1 were calculated, by considering the initial and final point of each test over the duration of the experimental run (i.e. t = 0 to t = 120–144 h). Three subsequent SBR experimental run (t = 41, 82, 120 days) were averaged together to summarize the values obtained along the SBR cultivation period. Maximum values indicate the maximum single data point measured during each individual SBR test. For the CR operations, samples were taken for analysis three times per week over a period of 90 days (n = 35). The average values reported in Table 1 show the average of the total amount of samples taken.

Under SBR conditions, an average volumetric productivities of 0.078 ± 0.012 g CDW/L·h was achieved. The latter value increased about 5-fold under CR configurations, reaching an average of 0.375 ± 0.015 g CDW/L·h. Biomass yields in terms of g CDW/g COD-H₂ increased from 0.073 ± 0.007 to 0.280 ± 0.010 g CDW/g COD-H₂, when changing from a SBR to a continuous operation mode. In the same way, CO₂-based yield increased from the minimum of 0.153 ± 0.023 g CDW-C/g CO₂-C observed during SBR cultivation to the maximum of 0.427 ± 0.013 g CDW-C/g CO₂-C. Hydrogen gas was also converted more efficiently when the reactor operated continuously, with an increase of 16% compared to SBR operations, reaching 81 ± 2%. Maximum values observed under CR were almost double than observed under SBR conditions. A different trend was observed for the nitrogen upgrade efficiency. SBR conditions allowed the complete conversion of the ammonium nitrogen supplied into MP, which reached an average of 65 ± 5% of the microbial biomass CDW. Under CR operation, instead, about 13% of the total mineral ammonium nitrogen supplied was still present in dissolved form in the CR effluent, whereas the average protein content of the produced biomass was 71 ± 5% (%CDW).

3.2. Microbial community analysis

In order to assess the composition of the microbial community, DNA samples from the SBR (after 120 days of operations) and from the CR configurations (after 20 days of operations) were analyzed by means of 16S rRNA Illumina sequencing.

The enriched microbial community cultivated under SBR conditions was characterized by a rather high diversity. Amongst the 12 genera identified, *Ancylobacter* (Morita, 1999), *Xanthobacter* (Wilde and Schlegel, 1982) and *Hydrogenophaga* (Willems et al., 1989) have

been already documented as able to carry out autotrophic oxyhydrogen metabolism. The latter constitute less than one-third of the quantitative genera distribution of the microbial community. No direct evidence of aerobic hydrogen oxidation is available in literature for the other genera present. Notably, the microbial community was dominated (one-third of the whole quantitative genera distribution) by *Bdellovibrio*, a genus of the class of *Deltaproteobacteria* encompassing predatory bacteria able to invade and lyse various other Gram-negative bacteria (Rendulic et al., 2004). The remaining genera detected (about 30% relative abundance) were mainly composed by the classes of *Falvobacteria* and *Sphingobacteria*, known as aerobic chemoorganotrophic bacteria (Vandamme et al., 1994; Yabuuchi et al., 1983).

Following the SBR cultivation period, the effect of the first 20 days of CR operations on the microbial community was investigated by means of a second 16S rRNA Illumina sequencing analysis. As shown in Fig. 1 b, the simple implementation of high rate (D = 0.1 h⁻¹) continuous reactor operations led to a remarkable selection within the microbial community, with almost 97% of the total community composed by a single genus: *Sulfuricurvum*. Almost 80% of the remaining 3% was composed by only other two genera: *Gammaproteobacteria* (*Thermomonas*) and *Flavobacteriia* (*Chryseobacterium*).

The DNA sample used for the 16S rRNA Illumina sequencing analysis of the CR was subsequently analyzed by means of 16S rRNA sequencing, together with a second sample taken after 90 days of continuous CR operations. The latter was done in order to confirm the stability of the microbial community composition and to gain more in depth information on the dominating *Sulfuricurvum* genus. For both samples the analysis indicated similarities at the level of 98 and 99% to *Sulfuricurvum kujiense* strains YK-2, YK-3 and YK-4, as well as to other uncultured *Epsilonproteobacteria* when compared using NCBI BLAST under default settings (Han et al., 2012).

3.3. Protein and amino acid profile

The bacterial biomass grown under CR configurations was harvested at day 90 (i.e. at the end of the CR cultivation period) and analyzed for crude protein content as well as for essential amino acids composition.

Fig. 2 compares the results obtained in this study with reference protein feed additives such as fishmeal, soybean meal and bacterial meal. The latter is a MP product obtained from methane oxidizing bacteria (*Methylococcus capsulatus* grown in association with other heterotrophic bacteria) already produced at pilot scale and tested in several feed trials involving monogastric animals as well as aquaculture species, for which the EU already approved the use in

Table 1
Parameters of HOB cultivation obtained under SBR tests (averaged over three different sequencing batch tests) and CR operations (over 90 days of continuous operations). Maximum values were calculated for each batch for the data points which maximized volumetric productivity and biomass yield, whereas average values were calculated over the whole period.

Parameter		Sequence batch reactor	Continuous reactor
Volumetric productivity (g CDW/L·h)	Average	0.078 ± 0.012	0.375 ± 0.015
	Maximum	0.187 ± 0.045	0.406
Y _{H₂} (g CDW/g COD-H ₂)	Average	0.073 ± 0.007	0.280 ± 0.010
	Maximum	0.157 ± 0.037	0.290
Y _{CO₂} (g CDW-C/g CO ₂ -C)	Average	0.153 ± 0.023	0.427 ± 0.013
	Maximum	0.246 ± 0.058	0.456
H ₂ gas conversion efficiency	Average	65% ± 4%	81% ± 2%
	Maximum	71% ± 3%	87%
N upgrade efficiency	Average	100%	87% ± 4%
	Maximum	100%	97%
Protein content (%CDW)	Average	66% ± 5%	71% ± 5%
	Maximum	73%	76%




Fig. 1. Phylogenetic composition of the HOB microbiome during SBR (a) and CR (b) operations, assessed by 16S rRNA Illumina sequencing. The timeline indicates the duration of each phase: SBR and CR, and when DNA samples were processed for 16S rRNA Illumina and Sanger sequencing. The central graph resumes the percentage of each class within the microbial community. Each class is then characterize in terms of genera composition in the external graphs (a, b) or within brackets (b).

animal nutrition (Øverland et al., 2010). Fishmeal and soybean meal were chosen as a reference for animal and vegetable protein, respectively. Bacterial meal allows to benchmark the MP produced in this study with another know similar product (i.e. already tested and legally approved MP).

As demonstrated in Fig. 2, the crude protein content of 71% of the *Sulfuricurvum* spp. microbial culture is slightly higher than bacterial meal (68%) and fishmeal (66%) and substantially higher than the average crude protein content of soybean meal (45%).

A similar trend can be observed in Fig. 3 for the amino acid profile. The profile for the *Sulfuricurvum* spp. microbial culture was comparable to that of bacterial meal and fishmeal and systematically better (at the exception of Arginine) than the one of soybean meal.

4. Discussion

4.1. Sequenced batch reactor

Following the enrichment, the SBR operations confirmed that the microbial culture effectively oxidized hydrogen coupled with assimilation of carbon dioxide and mineral nitrogen (i.e. ammonium nitrogen) into cell biomass. Consistent biomass growth was observed, allowing to operate the SBR at a SRT of about 6 days. Also, the $\text{NH}_4\text{-N}$ fed at the beginning of each SBR test was completely (100%) converted into organic nitrogen for microbial biomass build up. Nevertheless, the average performances observed in terms of volumetric productivities and biomass yield on hydrogen were far




Fig. 2. Crude protein content on CDW basis of the microbial biomass produced under CR configuration by the *Sulfuricurvum* spp. dominated culture (this study) compared with other microbial protein (bacterial meal), animal protein (fishmeal) and vegetable protein (soybean meal) (Øverland et al., 2010).




Fig. 3. Essential amino acids profile of the microbial biomass produced under CR configuration by the *Sulfuricurvum* spp. dominated culture (blue) (this study) compared with bacterial meal (yellow), fishmeal (grey) and soybean meal (green) as reported from Øverland et al. (2010). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

from being optimal. More specifically, the mixed culture grown under SBR conditions gave average biomass yields and productivities lower than values previously reported using specific bacterial strains (see Table 2).

The average volumetric productivity of 0.08 g CDW/L·h observed under SBR conditions, was 28.5, 9.1, 3.5, 6.4 and 3.3 times lower than the values reported for autotrophic growth of *Alcaligenes eutrophus*, *Alcaligenes eutrophus* ATCC 17697^T, *Ideonella* sp. O-1 and *Pseudomonas hydrogenovora*, respectively. Equally, the biomass yield on hydrogen gas was 3.8, 2.7 and 2.2 times lower than values reported for *Alcaligenes eutrophus* ATCC 17697^T, *Ideonella* sp. O-1 and *Pseudomonas hydrogenovora* grown under batch conditions.

The analysis of the community composition revealed a

surprising fractionation of the HOB enriched community into three distinct compartments: autotrophic HOB, heterotrophic bacteria and predatory bacteria, each sharing about 1/3 of the relative abundance of the overall community. The association between primary producers (autotrophic bacteria) and secondary consumers (heterotrophic bacteria) has already been documented in full scale MP production as well as reported and investigated in recent scientific studies (Aas et al., 2006; Ho et al., 2014). In the context of MP production, a clear example is represented by a methylotrophic bacterium (*Methylococcus capsulatus*) cultured in association with other heterotrophic bacteria. Such microbial fermentation is used in pilot-scale bioconversion of natural gas into MP (bacterial meal), eventually used as high-quality feed in aquaculture (Aas et al., 2006; Marit Berge et al., 2005). The coexistence of different microbial species offers benefits such as the removal of inhibiting byproducts or cell lysates, as well as the regulation of oxygen level (Ho et al., 2014; Strong et al., 2015).

Quite unexpected was the 33% relative abundance of *Bdellovibrio* spp., by far the most abundant genus dominating the mixed culture after 120 days of continuous SBR operations. The fact that such genus comprises predatory bacteria thriving on invasion and consumption of other Gram-negative bacteria (Rendulic et al., 2004) offers a reasonable yet remarkable explanation for the low performances of the HOB enriched community observed under SBR conditions. It is likely that the predatory activity of *Bdellovibrio* spp. imposed a major stress on the primary producers HOB, which were actively oxidizing hydrogen and fixing carbon dioxide into new microbial biomass then partly lysed by predatory activity. The lysed biomass might have also served as growth substrate for heterotrophic bacteria (Van Loosdrecht and Henze, 1999), in fact occupying the remaining 1/3 of the microbial community.

Table 2

Comparison of results obtained in this study for SBR and CR grown cultures with data from literature on single HOB strains.

Microbial culture/Strains	Substrate	Culture method	Biomass productivity (g CDW/L·h)	Biomass yield (g CDW/g COD-H ₂)	Reference
<i>Alcaligenes eutrophus</i>	H ₂ /O ₂ /CO ₂	Batch	2.28	–	(Tanaka et al., 1995)
<i>Alcaligenes eutrophus</i> ATCC 17697 ^T	H ₂ /O ₂ /CO ₂	Batch	0.71	0.28	(Ishizaki and Tanaka, 1990)
<i>Ideonella</i> sp. O-1	H ₂ /O ₂ /CO ₂	Batch	0.27	0.20	(Tanaka et al., 2011)
<i>Pseudomonas hydrogenovora</i>	H ₂ /O ₂ /CO ₂	Batch	0.50	0.16	(Goto et al., 1977)
Mixed culture (SBR)	H ₂ /Air/CO ₂	Batch	0.08	0.07	This study (average values)
<i>Alcaligenes eutrophus</i> ATCC17697	H ₂ /O ₂ /CO ₂	Continuous	0.40	0.29	(Morinaga et al., 1978)
<i>Alcaligenes hydrogenophilus</i>	H ₂ /O ₂ /CO ₂	Continuous	0.33	0.23	(Miura et al., 1982)
<i>Cupriavidus eutrophus</i> B-10646	H ₂ /O ₂ /CO ₂	Continuous	–	0.14	(Volova et al., 2013)
<i>Sulfuricurvum</i> spp. (CR)	H ₂ /Air/CO ₂	Continuous	0.38	0.28	This study (average values)

The high metabolic diversity and the low performances characterizing the microbial community under SBR condition can be also explained by speculating over the degrees of freedom of the biological systems in terms of growth rate and substrates concentrations, i.e. from a Monod-like point of view. Under SBR conditions, low constraints were imposed to the specific growth rate of the different bacteria present, which were therefore able to coexist in the same biological context. Also, the depletion of nutrients as well as the varying concentration of gasses as affected by the changing microbial activity over the batch culture (i.e. lag, log and decay phase), resulted in continuously changing growth conditions, potentially favoring different bacteria over time (see Fig. 4a).

In view of the sub-optimal performances, the SBR culture system did not seem to offer the best solution between process efficiency and stability. Moreover, the diverse microbial community would be difficult to control in terms of constancy of composition, and the presence of different bacterial strains of uncertain nutritional composition would affect the quality of the HOB microbiome as such for MP production for feed and food purposes.

4.2. Continuous reactor

The continuous operation at a dilution rate of 0.1 h^{-1} allowed to select for the evolvement of a more performing microbial culture in terms of biomass yields and volumetric productivities. Indeed the CR culture system selected for bacteria able to implement maximum substrate conversion at the specific growth rate imposed by the dilution D at which the bioreactor is operated (Goldberg, 1985). Thus microorganism having high specific growth rate can outcompete others not able to cope with the dynamics of the system. Such configuration can be summarized in the following two conditions:

- 1) $\mu \geq D = 0.1 \text{ h}^{-1}$: Continuous growth
- 2) $\mu < D = 0.1 \text{ h}^{-1}$: Wash-out

Fig. 4 b offers a virtual example of how the CR reactor impacted on the initial diverse community. The dilution rate of 0.1 h^{-1} required a corresponding specific growth rate of the same value. Moreover, the constant supply of nutrients and substrates to a biological system growing in steady conditions allowed to set a quite specific environment able to naturally select for the more adaptive and fast growing bacteria. In other terms, only bacteria possessing a specific growth rate higher than the dilution rate imposed, as well good affinities with the substrates provided were not washed-out.

As revealed by the microbial community analyses, within three weeks of operation the high dilution rate resulted in the selection of a highly specific microbial culture, dominated for more than 96% by *Sulfuricurvum* spp. The latter genus is known as encompassing a specific type of bacteria predominantly active towards sulfur oxidation in crude oil deposits (Han et al., 2012). *Sulfuricurvum kujiense* YK-1^T was first isolated from oil sands and characterized as a facultative anaerobic sulfur oxidizing bacteria (sulfide, elemental sulfur and thiosulfate) also able to use hydrogen as electron donor. Electron acceptors were described to be nitrate and oxygen under anaerobic and aerobic conditions, respectively. Aerobic growth though was limited to microaerophilic ranges (with maximum 1% in the headspace) (Kodama and Watanabe, 2004). Three other strains of *Sulfuricurvum kujiense* were already reported, but only strain YK-1^T was cultured and characterized in its whole genome (Han et al., 2012). Recently, the complete genome of *Candidatus Sulfuricurvum* sp. RIFRC-1 was assembled de novo from an aquifer-derived metagenome, confirming the importance and the link between sulfur and hydrogen metabolism in terrestrial subsurface environments. The latter finding also points out how more strains of the *Sulfuricurvum* spp. genus might still be discovered and characterized.

The *Sulfuricurvum* spp. dominating the culture studied in the present work is closely related to *Sulfuricurvum kujiense*, yet the exact identity of the strain is still unclear. The mineral medium used to culture the HOB both under sequenced batch and continuous reactor conditions did not contain any reduced sulfur compound such as sulfide, elemental sulfur or thiosulfate. Only oxidized sulfur under the form of sulfate ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.5 g/L)) was supplied to the reactor with the influent mineral medium. The development of the high enrichment of *Sulfuricurvum* spp. without any available reduced sulfur substrate points out that the most plausible metabolism for biomass growth within the biological system was hydrogen as electron donor and oxygen as electron acceptor. Therefore, when analyzing the conditions under which hydrogen oxidation was carried out, the fact that partial pressures of O_2 in the headspace of the CR were constantly higher than 1%, reaching 5–6% for long periods, constitutes a first important difference with the strain YK-1^T as characterized by Kodama and Watanabe (2004). The latter observation is supported by the fact that dissolved oxygen levels between 0.5 and 1.2 mg O_2/L could be measured in the effluent of the continuous reactor. Although the abovementioned physiological characterization reported the use of H_2 as electron donor in combination with microaerophilic O_2 concentrations, it did not identify the possibility of exploiting such bacterium for high rate autotrophic hydrogen oxidation, as experimentally





Fig. 4. Hypothetic resume of growth rates depending on substrates concentrations in SBR (a) and CR (b) conditions. (a) SBR culture conditions do not impose a strong growth rate to the system and substrate levels vary over the growth period, hence bacteria with different specific growth rates and affinities can freely co-evolve. (b) CR conditions are characterized by a strong dilution rate, which imposes a specific growth rate to the overall biological system. In the CR configuration substrates concentrations are well defined and become constant over time. Therefore only bacteria able to cope with the imposed dilution rate and having high affinity with the set of substrates concentrations can evolve in the system.

demonstrated in this study has not been described before.

As reported in Table 2, the cultured *Sulfuricurvum* spp. displayed biomass yields and volumetric productivities comparable to the ones reported for *Alcaligenes eutrophus* ATCC17697 from Morinaga et al. (1978), outscoring the values available in other studies for continuous cultures of *Alcaligenes hydrogenophilus* and *Cupriavidus eutrophus* B-10646 (Miura et al., 1982; Volova et al., 2013). *Alcaligenes eutrophus*, currently known as *Cupriavidus necator* can be also regarded as reference for a possible overall stoichiometry of carbon dioxide and ammonium nitrogen assimilation into bacterial biomass within the biological system driven by hydrogen oxidation (Ishizaki and Tanaka, 1990):



The molar ratios which can be calculated from the stoichiometry are in fact close to the gas ratios used in the present experiment. More specifically, from the stoichiometry a ratio of 3.4 can be calculated between H_2 and O_2 , which is comparable to the ratio of 3.2 at which the enrichment of *Sulfuricurvum* spp. was obtained. Similarly, the ratio of 4.8 between H_2 and CO_2 used in the present investigation was similar to the 5.2 obtained from the stoichiometry of *Cupriavidus necator*.

The fact that the culture dominated by *Sulfuricurvum* spp. matched efficiencies in terms of biomass yield and volumetric productivities of other well-known HOB strains while being fed with gas mixtures suitable for HOB growth, represents an interesting and novel finding and holds the potential to expand the biotech applications of autotrophic hydrogen oxidation to unexplored bacteria. Further research is warranted to investigate its potential in more detail.

Interestingly, the microbial composition was stable over the course of the experiments (90 days) and dominated by the same genus (see Fig. 1). This finding is important as this implies that the fermentation process can be easily managed without cumbersome sterility precautions (e.g. media autoclaving, gas filtering). The latter feature can be of interest in allowing the direct upgrade of used resources such as carbon dioxide and ammonia gas recovered e.g. from biogas and anaerobic digestate, respectively (Matassa et al., 2015b), without requiring strict subsequent axenic processing conditions and related operational costs. Further research is required to understand how such operational setting is resistant to external invasion and destabilization. Indeed, the latter can have biotechnological applications which go beyond the aim of this study.

In relation to the other bacteria coexisting with the *Sulfuricurvum* spp., the spectrum was composed by heterotrophic bacteria pertaining to the classes of *Gammaproteobacteria* (*Thermomonas*) and *Flavobacteriia* (*Chryseobacterium*). It is therefore likely that under high rate CR reactor configuration, these bacteria were benefitting from organic metabolites from the HOB, in this case *Sulfuricurvum* spp. Yet, this equilibrium achieved under CR conditions was totally different from the almost equal relative abundance between HOB and heterotrophs under SBR conditions. The magnitude of the residual heterotrophic niche (in terms of relative abundance) might be indeed depending on the growth conditions, and more specifically on the dilution rate imposed to the system. Further research efforts might aim at establishing whether or not such niche would be completely diminished at higher dilution rates, not allowing the secondary heterotrophic consumers to take advantage of the primary autotrophic carbon fixation activity.

4.3. Nitrogen assimilation efficiency, protein and amino acids profile

In terms of nitrogen assimilation and conversion efficiency, the system operated in batch-mode was able to convert 100% of NH_4-N nitrogen into MP at 66% or more protein content on CDW basis. In case of the CR configuration, the N-usage efficiency was lower, in the order of 87% on CDW basis. The aim of the high rate CR operation was to attain maximum biomass growth and MP accumulation, avoiding nutrient limitation. As result, nitrogen was added in a slight excess with some nitrogen was still present (unused) in the effluent of the reactor. It is likely that higher efficiencies could be obtained imposing more carefully N limiting conditions and varying the initial nitrogen loading rate.

The biomass produced under constant CR operation revealed a high protein content of more than 70%. The latter is in agreement with other studies on HOB for MP production (Volova and Barashkov, 2010), and confirms that *Sulfuricurvum* spp. might be suitable as a MP producing bacterium. The overall protein content is higher than the 68% reported for bacterial meal as well as than the one of fishmeal, regarded as high-quality additive in nutrition and also than the one of soybean meal, the reference vegetable protein for livestock. In the same way, the amino acids profile of the produced MP revealed a close compatibility to the one of bacterial meal as well as fishmeal, outscoring the one of soybean meal. Bacterial meal, as already produced from natural gas could also be used to directly upgrade the biogas produced from anaerobic digestion of sewage into MP. As discussed in a recent review (Matassa et al., 2015b), more than being self-excluding the hydrogen and the methane platforms can be seen as complementary, depending on the availability of each resource on-site. Like for bacterial meal, which already received positive feedback from feed trials, preliminary in vitro tests on the nutritional digestibility of our MP were also positive (data not shown). Clearly, the findings obtained in the study need to be complemented by detailed animal studies in which aspects of long-term gastro-intestinal uptake and putative nutritional side effects are scrutinized. However, the current findings clearly show the potential of using the produced MP as high-quality feed/food additive, offering a valid alternative to the high land, water, nutrients and carbon footprint of conventional vegetable protein production (Walsh et al., 2015). If this would be done by upgrading nitrogen recovered from used water the benefits in terms of avoided N losses and emissions could be even higher (Matassa et al., 2015a).

5. Conclusions

In this study, we aimed at assessing the potentialities of autotrophic hydrogen oxidation to recover and upgrade of resources under different operating conditions. The evolution of HOB from a generic mixed microbial community under different operating conditions allowed to reveal interesting and novel aspects, with potential for application in industrial contexts. The key findings are:

- Under SBR conditions the enriched mixed culture revealed the coexistence of a diversity of microbial actuators
- Under high rate CR culture conditions the microbiome narrowed down to *Sulfuricurvum* spp. dominated culture which was both stable and highly productive
- Mineral nitrogen and carbon dioxide were directly upgraded into microbial biomass, rich in protein, by using hydrogen and oxygen with high efficiency under CR culture conditions
- The nutritional properties of the produced MP are comparable to the high-quality fishmeal and surpass those of vegetable soybean meal.

Microbial biosynthesis of useful commodities from carbon dioxide is amongst the most challenging yet promising routes of the future bioeconomy. The exploration of renewable energy generation combined with technology advances in hydrogen production might enable on-site recovery and upgrading of valuable resources by means of HOB, produced under appropriate microbial resource management (MRM) conditions (Verstraete, 2015, 2007).

Acknowledgements

The authors express their gratitude to Elham Ehsani for her help with the DNA preparation for molecular analyses and to Esther Moens for her contribution to the bioreactor follow-up. Cindy Duysburgh is also gratefully acknowledged for her work on the preliminary tests on digestibility. The presented work was made possible by the MERMAID project “Microbial Resource Management and Engineering for Closing the Urban Water Cycle” a Marie Skłodowska-Curie Initial Training Network, financed by the FP7 of the European Commission under Grant number 607492 and by the Geconcerteerde Onderzoeksactie (GOA) of Ghent University (BOF15/GOA/006). The authors also acknowledge the Ghent University Multidisciplinary Partnership-Biotechnology for a sustainable economy (01 MRA 510 W) for supporting the presented work. The senior author also acknowledges the support of the Dutch watercycle research institute KWR.

References

- APHA, AWWA, WEF, 1992. Method 4500–NO₂: Standard Methods for the Examination of Water and Wastewater.
- Aas, T.S., Grisdale-Helland, B., Terjesen, B.F., Helland, S.J., 2006. Improved growth and nutrient utilisation in Atlantic salmon (*Salmo salar*) fed diets containing a bacterial protein meal. *Aquaculture* 259, 365–376. <http://dx.doi.org/10.1016/j.aquaculture.2006.05.032>.
- Agll, A.A.A., Hamad, Y.M., Hamad, T.A., Thomas, M., Bapat, S., Martin, K.B., Sheffield, J.W., 2013. Study of a molten carbonate fuel cell combined heat, hydrogen and power system: energy analysis. *Appl. Therm. Eng.* 59, 634–638. <http://dx.doi.org/10.1016/j.applthermaleng.2013.06.030>.
- Anupama, Ravindra, P., 2000. Value-added food: single cell protein. *Biotechnol. Adv.* 18, 459–479.
- Ball, M., Weeda, M., 2015. The hydrogen economy – vision or reality? This paper is also published as chapter 11 “the hydrogen economy – vision or reality?” in compendium of hydrogen energy volume 4: hydrogen use, safety and the hydrogen economy, edited by Michael Ball. Angelo Basi. *Int. J. Hydrogen Energy* 40, 7903–7919. <http://dx.doi.org/10.1016/j.ijhydene.2015.04.032>.
- Bartsev, S.I., Gitelson, J.I., Lisovsky, G.M., Mezhevikin, V.V., Okhonin, V.A., 1996. Perspectives of different type biological life support systems (BLSS) usage in space missions. *Acta Astronaut.* 39, 617–622. [http://dx.doi.org/10.1016/S0094-5765\(97\)00012-X](http://dx.doi.org/10.1016/S0094-5765(97)00012-X).
- Defoirdt, T., Boon, N., Sorgeloos, P., Verstraete, W., Bossier, P., 2007. Alternatives to antibiotics to control bacterial infections: luminescent vibriosis in aquaculture as an example. *Trends Biotechnol.* 25, 472–479. <http://dx.doi.org/10.1016/j.tibtech.2007.08.001>.
- Elsler, J.J., Fagan, W.F., Denno, R.F., Dobberfuhr, D.R., Folarin, A., Huberty, A., Interlandi, S., Kilham, S.S., McCauley, E., Schulz, K.L., Siemann, E.H., Sterner, R.W., 2000. Nutritional constraints in terrestrial and freshwater food webs. *Nature* 408, 578–580.
- Goldberg, I., 1985. *Biotechnology Monographs, vol. 4*. Springer-Verlag, Heidelberg.
- Goto, E., Suzuki, K., Kodama, T., Minoda, Y., 1977. Improvement of initial and exponential growth of hydrogen bacteria, strain 9-Y. *Agric. Biol. Chem.* 41, 521–525. <http://dx.doi.org/10.1080/00021369.1977.10862529>.
- Hamad, T.A., Agll, A.A., Hamad, Y.M., Bapat, S., Thomas, M., Martin, K.B., Sheffield, J.W., 2014. Study of combined heat, hydrogen and power system based on a molten carbonate fuel cell fed by biogas produced by anaerobic digestion. *Energy Convers. Manag.* 81, 184–191. <http://dx.doi.org/10.1016/j.enconman.2014.02.036>.
- Han, C., Kotsyurbenko, O., Chertkov, O., Held, B., Lapidus, A., Nolan, M., Lucas, S., Hammond, N., Deshpande, S., Cheng, J.-F., Tapia, R., Goodwin, L.A., Pitluck, S., Liolios, K., Pagani, I., Ivanova, N., Mavromatis, K., Mikhailova, N., Pati, A., Chen, A., Palaniappan, K., Land, M., Hauser, L., Chang, Y.-J., Jeffries, C.D., Brambilla, E.-M., Rohde, M., Spring, S., Sikorski, J., Göker, M., Woyke, T., Bristow, J., Eisen, J.A., Markowitz, V., Hugenholtz, P., Kyrpides, N.C., Klenk, H.-P., Detter, J.C., 2012. Complete genome sequence of the sulfur compounds oxidizing chemolithoautotroph *Sulfuricurvum kujiense* type strain (YK-1(T)). *Stand. Genomic Sci.* 6, 94–103. <http://dx.doi.org/10.4056/signs.2456004>.
- Ho, A., De Roy, K., Thas, O., De Neve, J., Hoefman, S., Vandamme, P., Heylen, K., Boon, N., 2014. The more, the merrier: heterotroph richness stimulates methanotrophic activity. *ISME J.* 8, 1945–1948.
- Ishizaki, A., Tanaka, K., 1990. Batch culture of alkaligenes eutrophus ATCC 17697 T using recycled gas closed circuit culture system. *J. Ferment. Bioeng.* 69, 170–174.
- Kodama, Y., Watanabe, K., 2004. *Sulfuricurvum kujiense* gen. nov., sp. nov., a facultatively anaerobic, chemolithoautotrophic, sulfur-oxidizing bacterium isolated from an underground crude-oil storage cavity. *Int. J. Syst. Evol. Microbiol.* 54, 2297–2300. <http://dx.doi.org/10.1099/ijs.0.63243-0>.
- Lee, J.Z., Logan, A., Terry, S., Spear, J.R., 2015. Microbial response to single-cell protein production and brewery wastewater treatment. *Microb. Biotechnol.* 8, 65–76. <http://dx.doi.org/10.1111/1751-7915.12128>.
- Liu, R., Wu, P., Lang, L., Xu, C., Wang, Y., 2016. Free Fe 3+/Fe 2+ improved the biomass resource recovery and organic matter removal in Rhodospirillum rubrum sphaeroides purification of sewage. *Environ. Technol.* 37, 108–113. <http://dx.doi.org/10.1080/09593330.2015.1064168>.
- Marit Berge, G., Baevefjord, G., Skrede, A., Storebakken, T., 2005. Bacterial protein grown on natural gas as protein source in diets for Atlantic salmon, *Salmo salar*, in saltwater. *Aquaculture* 244, 233–240. <http://dx.doi.org/10.1016/j.aquaculture.2004.11.017>.
- Matassa, S., Batstone, D.J., Huelsen, T., Schnoor, J.L., Verstraete, W., 2015a. Can direct conversion of used nitrogen to new feed and protein help feed the world? *Environ. Sci. Technol.* 49, 5247–5254. <http://dx.doi.org/10.1021/es505432w>.
- Matassa, S., Boon, N., Verstraete, W., 2015b. Resource recovery from used water: the manufacturing abilities of hydrogen-oxidizing bacteria. *Water Res.* 68, 467–478. <http://dx.doi.org/10.1016/j.watres.2014.10.028>.
- Miura, Y., Okazaki, M., Ohi, K., Nishimura, T., Komemushi, S., 1982. Optimization of biomass productivity and substrate utility of a hydrogen bacterium. *Biotechnol. Bioeng.* 24, 1173–1182.
- Morinaga, Y., Yamanaka, S., Ishizaki, A., Hirose, Y., 1978. Growth characteristics and cell composition of alkaligenes eutrophus in chemostat culture. *Yasushi Morinaga, Shigeru Yamanaka, Ayaaki Ishizaki and Y Oshio Hirose the central research laboratories of Ajinomoto Co., Inc., Kawasaki, Japan support system. Agric. Biol. Chem.* 42, 439–444. <http://dx.doi.org/10.1080/00021369.1978.10862993>.
- Morita, R., 1999. Is H₂ the universal energy source for long-term survival? *Microb. Ecol.* 38, 307–320. <http://dx.doi.org/10.1007/s002489901002>.
- Ni, M., Leung, D.Y.C., Leung, M.K.H., Sumathy, K., 2006. An overview of hydrogen production from biomass. *Fuel Process. Technol.* 87, 461–472. <http://dx.doi.org/10.1016/j.fuproc.2005.11.003>.
- Øverland, M., Tauson, A.-H., Shearer, K., Skrede, A., 2010. Evaluation of methane-utilising bacteria products as feed ingredients for monogastric animals. *Arch. Anim. Nutr.* 64, 171–189. <http://dx.doi.org/10.1080/17450391003691534>.
- Parkin, A., Sargent, F., 2012. The hows and whys of aerobic H₂ metabolism. *Curr. Opin. Chem. Biol.* 16, 26–34. <http://dx.doi.org/10.1016/j.cbpa.2012.01.012>.
- Pohlmann, A., Fricke, W.F., Reinecke, F., Kusian, B., Liesegang, H., Cramm, R., Eitinger, T., Ewering, C., Pötter, M., Schwartz, E., Strittmatter, A., Voss, I., Gottschalk, G., Steinbüchel, A., Friedrich, B., Bowen, B., 2006. Genome sequence of the bioplastic-producing “Knallgas” bacterium *Ralstonia eutropha* H16. *Nat. Biotechnol.* 24, 1–6. <http://dx.doi.org/10.1038/nbt1244>.
- Rendulic, S., Jagtap, P., Rosinus, A., Eppinger, M., Baar, C., Lanz, C., Keller, H., Lambert, C., Evans, K.J., Goesmann, A., Meyer, F., Sockett, R.E., Schuster, S.C., 2004. A predator unmasked: life cycle of *Bdellovibrio bacteriovorus* from a genomic perspective. *Science* 303, 689–692. <http://dx.doi.org/10.1126/science.1093027> (80-).
- Repaske, R., Mayer, R., 1976. Dense autotrophic cultures of alkaligenes eutrophus. *Appl. Environ. Microbiol.* 32, 592–597.
- Salo-väänänen, P.P., Koivistoinen, P.E., 1996. Determination of protein in foods: comparison of net protein and crude protein (N × 6.25) values. *Food Chem.* 57, 27–31. [http://dx.doi.org/10.1016/0308-8146\(96\)00157-4](http://dx.doi.org/10.1016/0308-8146(96)00157-4).
- Strong, P., Xie, S., Clarke, W.P., 2015. Methane as a resource: can the methanotrophs add value? *Environ. Sci. Technol.* <http://dx.doi.org/10.1021/es504242n>, 1502271040000007.
- Tanaka, K., Ishizaki, A., Kanamaru, T., Kawano, T., 1995. Production of poly(D-3-hydroxybutyrate) from CO₂, H₂, and O₂ by high cell density autotrophic cultivation of *Alcaligenes eutrophus*. *Biotechnol. Bioenergy* 45, 268–275. <http://dx.doi.org/10.1002/bit.260450312>.
- Tanaka, K., Miyawaki, K., Yamaguchi, A., Khosravi-Darani, K., Matsusaki, H., 2011. Cell growth and P(3HB) accumulation from CO₂ of a carbon monoxide-tolerant hydrogen-oxidizing bacterium, *Ideonella* sp. O-1. *Appl. Microbiol. Biotechnol.* 92, 1161–1169. <http://dx.doi.org/10.1007/s00253-011-3420-2>.
- Van Loosdrecht, M., Henze, M., 1999. Maintenance, endogenous respiration, lysis, decay and predation. *Water Sci. Technol.* 39, 107–117. [http://dx.doi.org/10.1016/S0273-1223\(98\)00780-X](http://dx.doi.org/10.1016/S0273-1223(98)00780-X).
- Vandamme, P., Bernadet, J., Segers, P., Kersters, K., Holmes, B., 1994. New perspectives in the classification of the flavobacteria : description of *Chryseobacterium* gen. nov., *Bergeyella* gen. nov. *Int. J. Syst. Bacteriol.* 44, 827–831.
- Verstraete, W., 2015. The technological side of the microbiome. *Npj Biofilms Microbiomes* 1, 15001.
- Verstraete, W., 2007. Microbial ecology and environmental biotechnology. *ISME J.* 1, 4–8. <http://dx.doi.org/10.1038/ismej.2007.7>.
- Vilchez-Vargas, R., Geffers, R., Suárez-Díez, M., Conte, I., Waliczek, A., Kaser, V.S., Kralova, M., Junca, H., Pieper, D.H., 2013. Analysis of the microbial gene landscape and transcriptome for aromatic pollutants and alkane degradation using a novel internally calibrated microarray system. *Environ. Microbiol.* 15, 1016–1039. <http://dx.doi.org/10.1111/j.1462-2920.2012.02752.x>.

- Volova, T.G., Barashkov, V.A., 2010. Characteristics of proteins synthesized by hydrogen-oxidizing microorganisms. *Appl. Biochem. Microbiol.* 46, 574–579. <http://dx.doi.org/10.1134/S0003683810060037>.
- Volova, T.G., Kiselev, E.G., Shishatskaya, E.I., Zhila, N.O., Boyandin, A.N., Syrvacheva, D.A., Vinogradova, O.N., Kalacheva, G.S., Vasiliev, A.D., Peterson, I.V., 2013. Cell growth and accumulation of polyhydroxyalkanoates from CO₂ and H₂ of a hydrogen-oxidizing bacterium, *Cupriavidus eutrophus* B-10646. *Bio-resour. Technol.* 146, 215–222. <http://dx.doi.org/10.1016/j.biortech.2013.07.070>.
- Walsh, B.J., Rydzak, F., Palazzo, A., Kraxner, F., Herrero, M., Schenk, P.M., Ciais, P., Janssens, I.A., Peñuelas, J., Niederl-Schmidinger, A., Obersteiner, M., 2015. New feed sources key to ambitious climate targets. *Carbon Balance Manag.* 10, 1–8. <http://dx.doi.org/10.1186/s13021-015-0040-7>.
- Westlake, R., 1986. Large-scale continuous production of single cell protein. *Chem. Ing. Technol.* 58, 934–937. <http://dx.doi.org/10.1002/cite.330581203>.
- Wilde, E., Schlegel, H.G., 1982. Oxygen tolerance of strictly aerobic hydrogen-oxidizing bacteria. *Ant. Van Leeuwenhoek* 48, 131–143.
- Willems, A., Busse, J., Goor, M., Pot, B., Falsen, E., Jantzen, E., Hoste, B., Gillis, M., Kersters, K., Auling, G., De Ley, J., 1989. *Hydrogenophaga*, a new genus of hydrogen-oxidizing bacteria that includes *hydrogenophaga flava* comb. nov. (formerly *pseudomonas flava*), *hydrogenophaga palleronii* (formerly *pseudomonas palleronii*), *hydrogenophaga pseudoflava* (formerly *pseudomonas pseudoflava*). *Int. J. Syst. Bacteriol.* 39, 319–333. <http://dx.doi.org/10.1099/00207713-39-3-319>.
- Yabuuchi, E., Kaneko, T., Yano, I., Moss, wayne, c., Miyoshi, N., 1983. *Sphingobacterium* Gen. nov., *Sphingobacterium Spiritivorum* Comb. nov., *Sphingobacterium Multivorum* Comb. nov., *Sphingobacterium Mizutae* Sp. nov., and *Flavobacterium Indologenes* Sp. Nov. : Glucose-nonfermenting Gram-negative Rods in CDC Groups IIK-2 and IIb, 33, pp. 580–598.
- Yu, J., Dow, A., Pingali, S., 2013. The energy efficiency of carbon dioxide fixation by a hydrogen-oxidizing bacterium. *Int. J. Hydrogen Energy* 38, 8683–8690. <http://dx.doi.org/10.1016/j.ijhydene.2013.04.153>.